


Natamycin®

(natamycin
ophthalmic suspension)
5%
Sterile

DESCRIPTION:

NATACYN® (natamycin ophthalmic suspension) 5% is a sterile, antifungal drug for topical ophthalmic administration. Each mL of NATACYN® (natamycin ophthalmic suspension) contains: **Active:** natamycin 5% (50 mg). **Preservative:** benzalkonium chloride 0.02%. **Inactive:** sodium hydroxide and/or hydrochloric acid (neutralized to adjust the pH), purified water.

The active ingredient is represented by the chemical structure:



Established Name: Natamycin
Molecular Formula: C₂₃H₄₇NO₁₃
Molecular Weight: 665.73 g/mol

possesses *in vitro* activity against a variety of yeast and filamentous fungi, including *Candida*, *Aspergillus*, *Cephalosporium*, *Fusarium* and *Penicillium*. The mechanism of action appears to be through binding of the molecule to the sterol moiety of the fungal cell membrane. The polyene-sterol complex alters the permeability of the membrane to produce depletion of essential cellular constituents. Although the activity against fungi is dose-related, natamycin is predominantly fungicidal. Natamycin is not effective *in vitro* against gram-positive or gram-negative bacteria. Topical administration appears to produce effective concentrations of natamycin within the corneal stroma but not in intraocular fluid. Systemic absorption should not be expected following topical administration of NATACYN® (natamycin ophthalmic suspension) 5%. As with other polyene antibiotics, absorption from the gastrointestinal tract is very poor. Studies in rabbits receiving topical natamycin revealed no measurable compound in the aqueous humor or sera, but the sensitivity of the measurement was no greater than 2 mg/mL.

INDICATIONS AND USAGE:

NATACYN® (natamycin ophthalmic suspension) 5% is indicated for the treatment of fungal blepharitis, conjunctivitis, and keratitis caused by susceptible organisms including *Fusarium solani* keratitis. As in other forms of suppurative keratitis, initial and sustained therapy of fungal keratitis should be determined by the clinical diagnosis, laboratory diagnosis by smear and culture of corneal scrapings and drug response. Whenever possible the *in vitro* activity of natamycin against the responsible fungus should be determined. The effectiveness of natamycin as a single agent in fungal endophthalmitis has not been established.

CONTRAINDICATIONS:

NATACYN® (natamycin ophthalmic suspension) 5% is contraindicated in individuals with a history of hypersensitivity to any of its components.

PRECAUTIONS:

General.

FOR TOPICAL OPHTHALMIC USE ONLY — NOT FOR INJECTION.

Failure of improvement of keratitis following 7-10 days of administration of the drug suggests that the infection may be caused by a microorganism not susceptible to natamycin.

Continuation of therapy should be based on clinical re-evaluation and additional laboratory studies.

Adherence of the suspension to areas of epithelial ulceration or retention of the suspension in the fornices occurs regularly. Use only if the container is undamaged.

Information for Patients:

Do not touch dropper tip to any surface, as this may contaminate the suspension. Patients

9012800

Read
Direction
(6039)



Chemical Name: Stereoisomer of 22-[(3-amino-3,6-dideoxy-β-D-mannopyranosyl)oxy]-1,3,26-trihydroxy-12-methyl-10-oxo-6,11,28-trioxatricyclo[22.3.1.05,7] octacosane, 8,14,16,18,20-pentaene-25-carboxylic acid.

Other: Pimaricin

The pH range is 5.0-7.5.

CLINICAL PHARMACOLOGY:

Natamycin is a tetraene polyene antibiotic derived from *Streptomyces natalensis*. It

EQUITY
PHARMACEUTICALS
APPROVED 02.07.2021
BY C.SNYMAN

Watson,
Beatriz

Digitally signed by Watson,
Beatriz
DN: cn=Watson, Beatriz,
ou=PDF Internal Signing,
o=Alcon Vision, LLC
Date: 2020.08.14 10:40:12
-05'00'

9018200_40351500_p3_LFT_X-4.indd 1

13/08/2020 17:22

		Brewery House, The Maltings, Silvester Street Kingston-Upon-Hull HU1 3JG Tel: +44 (0) 1482 973000	
Live Text:	<input checked="" type="checkbox"/> Yes / <input type="checkbox"/> No / <input type="checkbox"/> Both	Production Site:	EyeVance Pharmaceuticals
WO:	87	ACC:	Lori Loy
Comp. Description:	Leaflet_NATACYN_LDMO 50MG/ML GLB 15ML_US	Printing Colours:	Black
Comp. Number New:	9018200/40351500	Technical Colours:	Cutting Guides Dimensions
Comp. Number Old:	9016511 US	Minimum Font Size:	8pt
Format/Dimension:	5" x 11-15/16"	Font Type:	Helvetica Neue LT Pro, Symbol
Tech. Drawing No.:	L11965	Proof Number:	3 13/08/2020 Antonia Stark
Braille:			
! PLEASE TURN OVERPRINTING ON !			

should be advised not to wear contact lenses if they have signs and symptoms of fungal blepharitis, conjunctivitis, and keratitis.

Carcinogenesis, Mutagenesis, Impairment of Fertility:

There have been no long term studies done using natamycin in animals to evaluate carcinogenesis, mutagenesis, or impairment of fertility.

Pregnancy:

Animal reproduction studies have not been conducted with natamycin. It is also not known whether natamycin can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. NATACYN® (natamycin ophthalmic suspension) 5% should be given to a pregnant woman only if clearly needed.

Nursing Mothers:

It is not known whether these drugs are excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when natamycin is administered to a nursing woman.

Pediatric Use:

Safety and effectiveness in pediatric patients have not been established.

Geriatric Use:

No overall differences in safety or effectiveness have been observed between elderly and younger patients.

ADVERSE REACTIONS:

The following events have been identified during post-marketing use of NATACYN® (natamycin ophthalmic suspension) 5% in clinical practice. Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made. The events, which have been chosen for inclusion due to their seriousness, frequency of reporting, possible causal connection to NATACYN® (natamycin ophthalmic suspension) 5%, or a combination of these factors include: allergic reaction, change in vision, chest pain, corneal opacity, dyspnea, eye discomfort, eye edema, eye hyperemia, eye irritation, eye pain, foreign body sensation, paresthesia, and tearing.

DOSAGE AND ADMINISTRATION:

SHAKE WELL BEFORE USING. The preferred initial dosage in fungal keratitis is one drop of NATACYN® (natamycin ophthalmic suspension) 5% instilled in the conjunctival sac at hourly or two-hourly intervals. The frequency of application can usually be reduced to one drop 6 to 8 times daily after the first 3 to 4 days. Therapy should generally be continued for 14 to 21 days or until there is resolution of active fungal keratitis. In many cases, it may be helpful to reduce the dosage gradually at 4 to 7 day intervals to assure that the replicating organism has been eliminated. Less frequent initial dosage (4 to 6 daily applications) may be sufficient in fungal blepharitis and conjunctivitis.

HOW SUPPLIED:

NATACYN® (natamycin ophthalmic suspension) 5% is a 15 mL fill packaged in a 15 mL amber glass bottle with a black closure. A flint glass dropper with a red plastic closure and a black rubber bulb are packaged separately in a clear plastic blister with Tyvek backing.

15 mL NDC 71776-005-15

STORAGE: Store between 2°C to 24°C (36°F-75°F). *Do not freeze.* Avoid exposure to light and excessive heat.

Rx Only

Distributed by:
Eyevance Pharmaceuticals, LLC
Fort Worth, TX 76102
Revised: 05/2020
40351500
9018200



9018200_40351500_p3_LFT_X-4.indd 2

13/08/2020 17:22

				Brewery House, The Millings, Silvester Street Kingston-Upon-Hull HU1 3JG, Tel: +44 (0) 1482 973000	
Live Text:	<input checked="" type="checkbox"/> Yes / <input type="checkbox"/> No / <input type="checkbox"/> Both	Production Site: Eyevance Pharmaceuticals			
WO:	87	ACC: Lori Loy			
Comp. Description: Leaflet_NATACYN_LDMO 50MG/ML GLB 15ML_US		Printing Colours: Black			
Comp. Number New: 9018200/40351500		Technical Colours: Cutting Guides Dimensions			
Comp. Number Old: 9016511 US					
Format/Dimension: 5" x 11-15/16"					
Tech. Drawing No.: L11965					
Minimum Font Size: 8pt					
Font Type: Helvetica Neue LT Pro, Symbol					
Proof Number: 3 13/08/2020 Antonia Stark					
Braille:					
! PLEASE TURN OVERPRINTING ON !					